

An empirical mass formula for μ and τ leptons and some remarks on trident production

T. Jacobsen
Department of Physics, University of Oslo
PB 1048 Blindern
N-0316 Oslo, Norway

February 2, 2008

Abstract

For integer values of its free parameter, an empirical formula reproduces fairly well the mass values of the μ and τ leptons as if they were excited states of the electron. Trident production might possibly be due to constituent collisions.

PACS: 14.60.Cd, 14.60.Ef, 14.60.Fg, 14.60.Hi

An empirical mass formula

Three charged leptons are known, the electron and the heavier μ and τ leptons [1]. Relations between their mass values have been proposed [2]. It is therefore interesting to note that the empirical formula

$$m(n) = 3^{n/4}(4\pi)^{1+n}m_e,$$

where m_e is the mass of the electron, gives

$$m(1) = 106.199 \text{ MeV}/c^2$$

which is 0.5 per cent off the mass of the μ -lepton [1], and

$$m(2) = 1756.35 \text{ MeV}/c^2$$

which is 1.2 per cent off the mass of the τ -lepton [1]. That this formula reproduces the mass values of μ and of τ fairly well for integer values of n as if n is a quantum number suggests that μ and τ are excited states of the electron. If so, the electron would have to be a complex system of some constituents, e.g. rishons/preons [3, 4, 5].

Remarks on trident production

The electron and the proton are both stable particles. While a proton has constituents, has an electron constituents ?

If so, and if they by analogy to the three quarks in a proton were two x with electric charge $-2e/3$ each, and one y with electric charge $+e/3$, where each x and y has lepton number $1/3$, then a system with one x and two y could correspond to a neutrino.

An $x\bar{x}$ system with lepton number zero could correspond to a boson. If x and $\bar{x}$ analogous to quarks may be “dressed” as proposed in Fig. 1, an $x\bar{x}$ system may decay to a e^-e^+ pair.

“Spectators” and “leading particles” are seen in pd reactions where a beam proton collides with the neutron in the deuteron with its proton as spectator [6], and in $\bar{p}p$ reactions with a leading meson due to a dressed spectator quark [7]. If e^- and e^+ have some structure, leading spectator constituents could by analogy be expected also in e.g. e^-e^+ collisions. If they by analogy to quarks may be “dressed”, they may be observed.

If one x of a colliding e^- is scattered and dressed to be an $x\bar{x}$ boson which decays to an e^-e^+ pair, with the remaining xy system as a spectator which is dressed to be an xy electron as proposed in Fig. 2, the result is trident production, i.e.

$$e^- \rightarrow e^-e^+e^-.$$

Trident production has been observed [8] and discussed from theoretical point of view [9]. Our Fig. 2 suggests an additional production model.

Summary

Our empirical mass formula and model for trident production suggest that electrons may have an inner structure and some constituents.




Fig. 1.

Dressing of $x\bar{x}$ to an e^-e^+ pair as proposed in the text.




Fig. 2.

Separation and dressing of the assumed constituents of an electron to an electron and an $x\bar{x}$ pair as proposed in the text.

References

- [1] Particle Data Booklet 2004, LBNL and CERN, and Phys. Lett. B **592**, 1(2004).
- [2] A.O.Barut, Phys. Rev. Lett. **42**, 1251 (1979). P. Caldirola, Lett. Nuovo Cim. **27**, 225 (1980). H. Terazawa and M. Yasue, Phys. Lett. B **307**, 383 (1993). Y.Koide, arXiv:hep-ph/0506247 v1, 2005.
- [3] H. Harari, Phys. Lett. B **86**, 83 (1979), and Scientific American **248**, 48 (1983).
- [4] M.A. Shupe, Phys. Lett. B **86**, 87 (1979).
- [5] J.-J.Dugne, S.Fredriksson, and J.Hansson, Europhys. Lett. **57**, 188 (2002). S.Fredriksson, arXiv:hep-ph/0309213 v2, Sep. 2003. A.J.Buchmann and M.L.Schmid, Phys. Rev. D **71**, 055002(2005), and references therein. A.Breakstone, e-Print Arhive : physics/0602118, 2006, and references therein.
- [6] V.Bakken and T.Jacobsen, Nuovo Cim. A **61**, 219 (1981), and references therein.
- [7] F.Balestra et al., Phys. Lett. B **217**, 43 (1989). F. Balestra et al., Phys. Scripta **43**, 9 (1991).
- [8] P.H.Fowler, D.H.Perkins and C.F.Powell, The Study of Elementary Particles by the Photographic Method, Pergamon Press, London 1959, and references therein.
- [9] G.Reading Henry, Phys. Rev. **154**, 1534 (1967), K.A.Thompson and P.Chen, SLAC PUB 7776 (1998), T.Adams et al., hep-ex/9811012, 9 Nov 1998, and references therein.